


azstarys® II
serdexmethylphenidate
and dexamethylphenidate
26.1/5.2mg • 39.2/7.8mg • 52.3/10.4mg capsules

ANNOUNCING THE NEW COPAY SAVINGS OFFER

Eligible patients **PAY \$0** for their first AZSTARYS® prescription*

\$25 for refills if their insurance covers AZSTARYS

\$50 for refills if their insurance does not cover AZSTARYS



This savings offer is available for a limited time

*Eligibility required. Restrictions may apply. See Terms and Conditions at [AZSTARYS.com](https://www.azstarys.com).

Eligible patients can register for their copay savings card at [AZSTARYS.com](https://www.azstarys.com) to get started



Scan the QR code or call **1-800-910-8432** to learn more about copay savings and support.

INDICATION

AZSTARYS is a central nervous system (CNS) stimulant indicated for the treatment of Attention Deficit Hyperactivity Disorder (ADHD) in patients 6 years and older.

IMPORTANT SAFETY INFORMATION

WARNING: ABUSE AND DEPENDENCE

- CNS stimulants, including AZSTARYS, other methylphenidate-containing products, and amphetamines, have a high potential for abuse and dependence. Assess the risk of abuse prior to prescribing, and monitor for signs of abuse and dependence while on therapy

Please see additional Important Safety Information on back page and Full Prescribing Information, including Boxed Warning, by visiting [AZSTARYS-pro.com](https://www.azstarys-pro.com).

IMPORTANT SAFETY INFORMATION (continued)

Contraindications

- Known hypersensitivity to serdexmethylphenidate, methylphenidate, or other product components. Bronchospasm, rash, and pruritus have occurred with AZSTARYS. Hypersensitivity reactions such as angioedema and anaphylactic reactions have occurred with other methylphenidate products.
- Concomitant treatment with a monoamine oxidase inhibitor (MAOI) or use of an MAOI within the preceding 14 days, because of the risk of hypertensive crisis.

Warnings and Precautions

- Sudden death has been reported in association with CNS stimulant treatment at recommended doses in pediatric patients with structural cardiac abnormalities or other serious heart problems. In adults, sudden death, stroke, and myocardial infarction have been reported at recommended doses. Avoid use in patients with known structural cardiac abnormalities, cardiomyopathy, serious heart arrhythmias, coronary artery disease, or other serious heart problems.
- CNS stimulants cause an increase in blood pressure and heart rate. Monitor all patients for hypertension and tachycardia.
- *Exacerbation of Pre-existing Psychosis:* May exacerbate symptoms of behavior disturbance and thought disorder in patients with a pre-existing psychotic disorder. *Induction of a Manic Episode in Patients with Bipolar Disorder:* May induce a mixed/manic episode in patients with bipolar disorder. Prior to initiating treatment, screen for risk factors for developing a manic episode (e.g., comorbid or history of depressive symptoms, or a family history of suicide, bipolar disorder, or depression). *New Psychotic or Manic Symptoms:* At recommended doses, may cause psychotic or manic symptoms (e.g., hallucinations, delusional thinking, or mania) in patients without a history of psychotic illness or mania. Discontinue if symptoms occur.
- Cases of painful and prolonged penile erections and priapism have been reported with methylphenidate products. Immediate medical attention should be sought if signs or symptoms of prolonged penile erections or priapism are observed.
- CNS stimulants, including AZSTARYS, are associated with peripheral vasculopathy, including Raynaud's phenomenon. Signs and symptoms are usually intermittent and mild; very rare sequelae include digital ulceration and/or soft tissue breakdown. Carefully observe patients during treatment for digital changes. Further evaluation may be required, including referral.
- CNS stimulants have been associated with weight loss and slowing of growth rate in pediatric patients. Monitor height and weight at appropriate intervals in pediatric patients. Treatment may need to be interrupted in children not growing or gaining weight as expected.

Adverse Reactions

- Based on accumulated data from other methylphenidate products, the most common (>5% and twice the rate of placebo) adverse reactions are appetite decreased, insomnia, nausea, vomiting, dyspepsia, abdominal pain, weight decreased, anxiety, dizziness, irritability, affect lability, tachycardia, and blood pressure increased.

Drug Interactions

- Adjust dosage of antihypertensive drug as needed. Monitor blood pressure.
- Avoid use of AZSTARYS on the day of surgery if halogenated anesthetics will be used.

Please see additional Important Safety Information on front page and Full Prescribing Information, including Boxed Warning, by visiting AZSTARYS-pro.com.

Corium

© 2022 Corium, Inc. All Rights Reserved.
COM-AZS-001694 (v 2.0) 05/22



azstarys®

serdexmethylphenidate
and dexamethylphenidate

26.1/5.2mg • 39.2/7.8mg • 52.3/10.4mg capsules